

The role of continuum and three-nucleon forces in neutron rich nuclei.

Gaute Hagen (ORNL)

Collaborators:

Morten Hjorth-Jensen (UiO/CMA)


Gustav Jansen (UT/ORNL)

Ruprecht Machleidt (UI)

Thomas Papenbrock (UT/ORNL)

TRIUMF Seminar

Vancouver, February 13 2013




Outline

1. Interactions from chiral EFT, open quantum systems and Coupled-Cluster theory
2. Evolution of shell structure in neutron rich calcium isotopes – Is ^{54}Ca a magic nucleus?
3. Role of continuum and three-nucleon forces in neutron rich oxygen and fluorine isotopes
4. Proton elastic scattering of ^{40}Ca using CC
5. Computing scattering observables from a finite Harmonic oscillator basis


Roadmap for Theory of Nuclei

Main goal :



To arrive at a comprehensive description of all nuclei and low-energy reactions from the basic interactions between the constituent nucleons




What are the relevant degrees of freedom?



Effective field theory provides us with a systematic link between quarks and gluons and nuclei.



Three-nucleon forces as in-medium corrections to nucleon-nucleon forces



Integrating over the third leg in infinite nuclear matter and derive density dependent corrections to the nucleon-nucleon interaction.


J. W. Holt N. Kaiser and W. Weise. Phys.Rev.C 79, 054331 (2009)
K. Hebeler and A. Schwenk (2010)

Our strategy: C_D is given by fit to triton half-life, we fix C_E and k_F from fit to binding energy in selected medium mass nuclei:

Schematic three-nucleon forces

Including the effects of 3NFs (approximation!)

[J.W. Holt, Kaiser, Weise, PRC 79, 054331 (2009); Hebeler & Schwenk, PRC 82, 014314 (2010)]




3NFs as in-medium effective two-nucleon forces

Integration of Fermi sea of symmetric nuclear matter: k_F

Parameters: For Oxygen we use $k_F=1.05 \text{ fm}^{-1}$, $c_E=0.71$, $c_D=-0.2$ from binding energies of $^{16,22}\text{O}$, for Calcium we use $k_F = 0.95 \text{ fm}^{-1}$, $c_E = 0.735$, $c_D = -0.2$ from binding energy of ^{40}Ca and ^{48}Ca (The parameters c_D , c_E differ from values proposed for light nuclei)




Physics of nuclei at the edges of stability



The Berggren completeness treats bound, resonant and scattering states on equal footing.

Has been successfully applied in the shell model in the complex energy plane to light nuclei. For a review see

N. Michel et al J. Phys. G 36, 013101 (2009).



Coupled-cluster method (in CCSD approximation)

Ansatz:

$$|\Psi\rangle = e^T |\Phi\rangle$$


$$T = T_1 + T_2 + \dots$$

$$T_1 = \sum_{ia} t_i^a a_a^\dagger a_i$$

$$T_2 = \sum_{ijab} t_{ij}^{ab} a_a^\dagger a_b^\dagger a_j a_i$$

- ☺ Scales gently (polynomial) with increasing problem size $\mathcal{O}(u^4)$.
- ☺ Truncation is the only approximation.
- ☺ Size extensive (error scales with A)
- ☹ Most efficient for doubly magic nuclei

Correlations are *exponentiated* 1p-1h and 2p-2h excitations. Part of np-nh excitations included!



Coupled cluster equations

$$E = \langle \Phi | \bar{H} | \Phi \rangle$$


$$0 = \langle \Phi_i^a | \bar{H} | \Phi \rangle$$

$$0 = \langle \Phi_{ij}^{ab} | \bar{H} | \Phi \rangle$$

Alternative view: CCSD generates similarity transformed Hamiltonian with no 1p-1h and no 2p-2h excitations.


$$\bar{H} \equiv e^{-T} H e^T = (H e^T)_c = \left(H + H T_1 + H T_2 + \frac{1}{2} H T_1^2 + \dots \right)_c$$

Evolution of shell structure in neutron rich Calcium



- How do shell closures and magic numbers evolve towards the dripline?
- Is the naïve shell model picture valid at the neutron dripline?

- 3NFs are responsible for shell closure in ^{48}Ca
 - Different models give conflicting result for shell closure in ^{54}Ca .
- J. D. Holt et al, J. Phys. G **39**, 085111 (2012)




Evolution of shell structure in neutron rich Calcium

Inversion of shell order in ^{60}Ca

S. M. Lenzi, F. Nowacki, A. Poves, and K. Sieja Phys. Rev. C 82, 054301 (2010)


- Inversion of d5/2 and g9/2 in ^{60}Ca .
- Bunching of levels pointing to no shell-closure.



Evolution of shell structure in neutron rich Calcium

- Relativistic mean-field show no shell gap in $60-70\text{Ca}$
- Bunching of single-particle orbitals
- large deformations and no shell closure


J. Meng et al, Phys. Rev. C 65, 041302(R) (2002)



How many protons and neutrons can be bound in a nucleus?

Literature: 5,000-12,000

Skyrme-DFT: $6,900 \pm 500_{\text{syst}}$




Description of observables and model-based extrapolation

- Systematic errors (due to incorrect assumptions/poor modeling)
- Statistical errors (optimization and numerical errors)

Erler et al., Nature 486, 509 (2012)

Calcium isotopes from chiral interactions



Main Features:


1. Total binding energies agree well with experimental masses.
2. Masses for $^{40-52}\text{Ca}$ are converged in 19 major shells.
3. ^{60}Ca is not magic
4. $^{61-62}\text{Ca}$ are located right at threshold.

See also:

Meng et al PRC 65, 041302 (2002), Lenzi et al PRC 82, 054301 (2010) and Erler et al, Nature 486, 509 (2012)

G. Hagen, M. Hjorth-Jensen, G. R. Jansen, R. Machleidt, T. Papenbrock, Phys. Rev. Lett. 109, 032502 (2012).

Is ^{54}Ca a magic nucleus? (Is $N=34$ a magic number?)



Main Features:


1. Good agreement between theory and experiment.
2. Shell closure in ^{48}Ca due to effects of 3NFs
3. Predict weak (sub-)shell closure in ^{54}Ca .

Measurement at RIKEN (Japan) agrees with theoretical prediction.

Hagen, Hjorth-Jensen, Jansen, Machleidt, T. Papenbrock, Phys. Rev. Lett. 109, 032502 (2012).

| ^{48}Ca | | | ^{52}Ca | | | ^{54}Ca | | |
|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|
| 2^+ | 4^+ | $4^+/2^+$ | 2^+ | 4^+ | $4^+/2^+$ | 2^+ | 4^+ | $4^+/2^+$ |
| CC 3.58 | 4.20 | 1.17 | 2.19 | 3.95 | 1.80 | 1.89 | 4.46 | 2.36 |
| Exp 3.83 | 4.50 | 1.17 | 2.56 | ? | ? | ? | ? | ? |

Spectra and shell evolution in Calcium isotopes



1. Inversion of the $9/2^+$ and $5/2^+$ resonant states in $^{53,55,61}\text{Ca}$
2. We find the ground state of ^{61}Ca to be $\frac{1}{2}^+$ located right at threshold.
3. A harmonic oscillator basis gives the naïve shell model ordering of states.

Continuum coupling is crucial!

| | ^{48}Ca | ^{52}Ca | ^{54}Ca |
|-------------------------------|------------------|------------------|------------------|
| $E_{2^+}(\text{CC})$ | 3.58 | 2.19 | 1.89 |
| $E_{2^+}(\text{Exp})$ | 3.83 | 2.56 | n.a. |
| $E_{4^+}/E_{2^+}(\text{CC})$ | 1.17 | 1.80 | 2.36 |
| $E_{4^+}/E_{2^+}(\text{Exp})$ | 1.17 | n.a. | n.a. |
| $S_n(\text{CC})$ | 9.45 | 6.59 | 4.59 |
| $S_n(\text{Exp})$ | 9.95 | 6.0* | 4.0† |




New penning trap measurement of masses of $^{51,52}\text{Ca}$
A. T. Gallant et al Phys. Rev. Lett. **109**, 032506 (2012)

| | ^{53}Ca | | ^{55}Ca | | ^{61}Ca | |
|---------|------------------|----------|------------------|----------|------------------|----------|
| J^π | Re[E] | Γ | Re[E] | Γ | Re[E] | Γ |
| $5/2^+$ | 1.99 | 1.97 | 1.63 | 1.33 | 1.14 | 0.62 |
| $9/2^+$ | 4.75 | 0.28 | 4.43 | 0.23 | 2.19 | 0.02 |

Is ^{28}O a bound nucleus?

Experimental situation

- “Last” stable oxygen isotope ^{24}O
- $^{25,26}\text{O}$ unstable (Hoffman et al 2008, Lunderberg et al 2012)
- ^{28}O not seen in experiments
- ^{31}F exists (adding on proton shifts drip line by 6 neutrons!?)



Continuum shell model with HBUSD interaction predict ^{28}O unbound. A. Volya and V. Zelevinsky PRL (2005)



Shell model (sd shell) with monopole corrections based on three-nucleon force predicts ^{24}O as last stable isotope of oxygen. [Otsuka, Suzuki, Holt, Schwenk, Akaishi, PRL (2010), arXiv:0908.2607]

Oxygen isotopes from chiral interactions



1. Effective 3NF place dripline at ^{25}O .
2. Odd-even staggering is well reproduced.
3. ^{26}O unbound by $\sim 100\text{keV}$,
**Lunderberg et al.,
Phys. Rev. Lett. 108
(2012) 142503**
4. ^{28}O unbound, with a width of $\sim 2\text{MeV}$

G. Hagen, M. Hjorth-Jensen, G. R. Jansen, R. Machleidt, T. Papenbrock, Phys. Rev. Lett. 108, 242501 (2012).

Chiral three-nucleon force at order N2LO. $k_f = 1.05\text{fm}^{-1}$,
 $C_D = 0.2$, $C_E = 0.71$ (k_f and c_E fitted to the binding Energy of ^{16}O and ^{22}O).




Resonances in neutron rich oxygen-24



C. R. Hoffman et al Phys. Rev. C **83**, 031303(R) (2011)

- Knockout reaction of ^{26}F reveal a resonance above the two-neutron threshold in ^{24}O
- No spin and parity assigned of this state
- A challenge for microscopic theory to address these states

Oxygen isotopes from chiral interactions



The effects of three-nucleon forces decompress the spectra and brings it in good agreement with experiment.

We find several states (4^+ , 3^+ , 2^+) near the observed peak at $\sim 7.5\text{MeV}$ in ^{24}O C. R. Hoffman et al Phys. Rev. C **83**, 031303 (2011)

| J^π | 2_1^+ | 1_1^+ | 4_1^+ | 3_1^+ | 2_2^+ | 1_2^+ |
|----------------|------------------------|------------------------|---------|---------|---------|---------|
| E_{CC} | 4.56 | 5.2 | 6.2 | 6.9 | 7.0 | 8.4 |
| E_{Exp} | 4.7(1) | 5.33(10) | | | | |
| Γ_{CC} | 0.03 | 0.04 | 0.005 | 0.01 | 0.04 | 0.56 |
| Γ_{Exp} | $0.05^{+0.21}_{-0.05}$ | $0.03^{+0.12}_{-0.03}$ | | | | |

Hagen, Hjorth-Jensen, Jansen, Machleidt, T. Papenbrock, Phys. Rev. Lett. **108**, 242501 (2012).



Computing open-shell Fluorine-26

$$(\bar{H} \hat{R}_\mu^{(A\pm 2)})_C |\Phi_0\rangle = \omega_\mu \hat{R}_\mu^{(A\pm 2)} |\Phi_0\rangle$$

$$\hat{R}^{(A+2)} = \frac{1}{2} \sum_{ba} r^{ab} a_a^\dagger a_b^\dagger + \frac{1}{6} \sum_{iabc} r_i^{abc} a_a^\dagger a_b^\dagger a_c^\dagger a_i + \dots$$

π -protons

ν -neutrons



Experimental spectra in ^{26}F compared with phenomenological USD shell-model calculations and coupled-cluster calculations

A. Lepailleur et al, accepted for publication in PRL(2012)



Elastic proton/neutron scattering on ^{40}Ca

The one-nucleon overlap function: $O_A^{A+1}(lj; kr) = \sum_n \left\langle A+1 \left\| \tilde{a}_{nlj}^\dagger \right\| A \right\rangle \phi_{nlj}(r)$.


Beyond the range of the nuclear interaction the overlap functions take the form:

$$O_A^{A+1}(lj; kr) = C_{lj} \frac{W_{-\eta, l+1/2}(kr)}{r}, \quad k = i\kappa$$

$$O_A^{A+1}(lj; kr) = C_{lj} [F_{\ell, \eta}(kr) - \tan \delta_l(k) G_{\ell, \eta}(kr)]$$




Elastic proton/neutron scattering on ^{40}Ca




Differential cross section for elastic proton scattering on ^{40}Ca .

Fair agreement between theory and experiment for low-energy scattering.

G. Hagen and N. Michel
Phys. Rev. C **86**, 021602(R) (2012).



What is the infrared cutoff in harmonic oscillator expansions?



Top: Exact wavefunction for a square well potential and the wavefunction from a HO basis expansion with $N=4$ and $\hbar\omega = 6\text{MeV}$. Bottom: Wavefunction imposing Dirichlet boundary conditions at L_0 , L'_0 , L_2

For energy/radii extrapolation in a finite HO basis
Furnstahl, Hagen, Papenbrock,
PRC 86, 031301 (2012)

A finite Harmonic oscillator basis expansion effectively imposes box boundary conditions at L . Candidates for the infrared cutoff L :

$$L_0 = \sqrt{2(N + 3/2)b}$$


$$L'_0 = L_0 + 0.54437b(L_0/b)^{-1/3}$$

$$L_2 = \sqrt{2(N + 3/2 + 2)b}$$

Correct IR cutoff

The energy with Dirichlet boundary conditions at L :

$$E(L) = E_\infty + Ae^{-2k_\infty L} + \mathcal{O}(e^{-4k_\infty L})$$



Solving a two-body problem exactly we can by inspection determine which infrared cutoff is favored.

Clearly L_2 is the right choice!

S. N. More et al, to be published (2013)

Numerical and analytical derivation of L_2


- For s-waves the lowest eigenvalue k_0 of k^2 in a box of size L is π/L
- The lowest eigenvalue of k^2 in a finite HO basis is to a good approximation given by π/L_2
- A finite HO basis imposes a Dirichlet boundary conditions approximately at L_2 , it can be shown to be exact for $N \gg 1$

| N | κ_{\min} | π/L_2 | π/L_0 |
|-----|-----------------|-----------|-----------|
| 0 | 1.2247 | 1.1874 | 1.8138 |
| 2 | 0.9586 | 0.9472 | 1.1874 |
| 4 | 0.8163 | 0.8112 | 0.9472 |
| 6 | 0.7236 | 0.7207 | 0.8112 |
| 8 | 0.6568 | 0.6551 | 0.7207 |
| 10 | 0.6058 | 0.6046 | 0.6551 |
| 12 | 0.5651 | 0.5642 | 0.6046 |
| 14 | 0.5316 | 0.5310 | 0.5642 |
| 16 | 0.5035 | 0.5031 | 0.5310 |
| 18 | 0.4795 | 0.4791 | 0.5031 |
| 20 | 0.4585 | 0.4582 | 0.4791 |

We can determine the exact box boundary L for a finite HO basis expansion, by diagonalizing k^2 and for each discrete k_i solve for

$$j_l(k_i L) = 0$$

For increasing HO basis size, the box boundary L approaches L_2 over a large range of energies.





Phaseshifts from a finite Harmonic Oscillator basis

- Diagonalize k^2 in a given finite localized basis such as the Harmonic Oscillator.
- Determine the infrared cutoff or box boundary L as a function of energy by solving the i 'th root of

$$j_l(k_i L) = 0$$

- Diagonalize the full Hamiltonian in the given finite basis and obtain the phase shifts from

$$\tan \delta(k_i) = \frac{j_l(k_i L)}{n_l(k_i L)}$$



Summary

1. Interactions from Chiral EFT probed in nuclei
2. CC calculations for oxygen and calcium with effects of 3NF and continuum give significant improvement in binding energy and spectra.
3. Predict weak sub-shell closure in ^{54}Ca .
4. Level ordering in the *gds* shell in neutron calcium is reversed compared to naïve shell model.
5. Predict spin and parity of newly observed resonance peak in ^{24}O .
6. Elastic proton scattering on medium mass nuclei from coupled-cluster theory
7. Phaseshifts from finite harmonic oscillator bases

Treatment of long-range Coulomb effects

We write the Coulomb interaction

$$V_{\text{Coul}} = U_{\text{Coul}}(r) + [V_{\text{Coul}} - U_{\text{Coul}}(r)]$$

Demanding

$$U_{\text{Coul}}(r) \longrightarrow (Z - 1)e^2/r \text{ for } r \rightarrow +\infty$$


The second term is short range and can be Expanded in Harmonic Oscillator basis. The first term contain the long range Coulomb part:

$$U_{\text{Coul}}(k, k') = \langle k | U_{\text{Coul}}(r) - \frac{(Z - 1)e^2}{r} | k' \rangle + \frac{(Z - 1)e^2}{\pi} Q_\ell \left(\frac{k^2 + k'^2}{2kk'} \right)$$


We diagonalize the one-body Schrödinger equation in momentum space using the off-diagonal method

N. Michel Phys. Rev. C 83, 034325 (2011)

| | | $s_{1/2}$ | | $d_{3/2}$ | | $d_{5/2}$ | |
|---------------|-------|-----------|----------|-----------|----------|-----------|----------|
| N_R | N_T | Re[E] | Γ | Re[E] | Γ | Re[E] | Γ |
| 5 | 15 | 1.1054 | 0.1446 | 5.0832 | 1.3519 | 1.4923 | 0.0038 |
| 5 | 20 | 1.1033 | 0.1483 | 5.0785 | 1.3525 | 1.4873 | 0.0079 |
| 10 | 25 | 1.0989 | 0.1360 | 5.0765 | 1.3525 | 1.4858 | 0.0093 |
| 10 | 30 | 1.0986 | 0.1366 | 5.0757 | 1.3529 | 1.4849 | 0.0103 |
| 15 | 40 | 1.0978 | 0.1351 | 5.0749 | 1.3531 | 1.4842 | 0.0111 |
| 15 | 50 | 1.0978 | 0.1353 | 5.0746 | 1.3533 | 1.4838 | 0.0114 |
| 20 | 60 | 1.0976 | 0.1349 | 5.0745 | 1.3533 | 1.4837 | 0.0116 |
| 30 | 70 | 1.0975 | 0.1346 | 5.0744 | 1.3534 | 1.4837 | 0.0117 |
| (Michel 2011) | | 1.0975 | 0.1346 | 5.0744 | 1.3535 | 1.4836 | 0.0119 |



Evolution of the $3/2^+$ and $1/2^+$ states in Potassium



1. We reproduce level inversion in ^{47}K and get $1/2^+$ as the ground state.
2. We predict $3/2^+$ for the ground state in $^{51,53,59}\text{K}$.

$1/2^+$ state with respect to $3/2^+$ state in ^{39}K and ^{47}K .

^{39}K

^{47}K

| J^π | E(CC) | E(Exp) | E(CC) | E(Exp) |
|---------|-------|--------|--------|--------|
| $3/2^+$ | 0.00 | 0.00 | 0.00 | 0.00 |
| $1/2^+$ | 4.097 | 2.52 | -0.636 | -0.36 |